

The Union Jack model: a quantum spin model with frustration on the square lattice

A. Collins, J. McEvoy, D. Robinson, C. J. Hamer and Zheng Weihong¹

¹*School of Physics, The University of New South Wales, Sydney, NSW 2052, Australia*

(Dated: February 2, 2008)

A new quantum spin model with frustration, the ‘Union Jack’ model on the square lattice, is analyzed using spin-wave theory. For small values of the frustrating coupling α , the system is Néel ordered as usual, while for large α the frustration is found to induce a canted phase. The possibility of an intermediate spin-liquid phase is discussed.

PACS numbers: PACS Indices: 05.30.-d, 75.10.-b, 75.10.Jm, 75.30.Ds, 75.30.Kz

(Submitted to Phys. Rev. B)

I. INTRODUCTION

Frustrated lattice spin models in two dimensions have attracted much discussion in recent years. They exhibit new and interesting phase structures and phase transitions; in particular, they may develop ‘spin liquid’ states, without long-range order [1]. It is also believed that frustrating interactions may play a role in the high-temperature superconducting cuprate materials. Primary examples are the anisotropic triangular lattice Heisenberg antiferromagnet (Fig. 1a), the square lattice $J_1 - J_2$ model (Fig. 1b), and the Shastry-Sutherland model (Fig. 1c). In this paper we discuss another member of this group, the ‘Union Jack’ model (Fig. 1d), which is another square lattice Heisenberg antiferromagnet with frustration, and might be expected to display some interesting properties.

The spin-1/2 $J_1 - J_2$ model on the square lattice [2] involves antiferromagnetic Heisenberg spin interactions with coupling J_1 between nearest neighbours, and coupling J_2 between diagonal next-nearest neighbours, as illustrated in Figure 1b). The Union Jack model has J_2 interactions on only half the diagonal bonds, in the pattern shown in Figure 1d), so that the lattice consists of two different site types, A and B, and the unit cell is 2x2 sites. Both models will exhibit quantum phase transitions as the coupling ratio $\alpha = J_2/J_1$ is varied.

In the $J_1 - J_2$ model at small α , the J_1 coupling is dominant and produces antiferromagnetic Néel ordering of the spins. At large α , the J_2 interaction is dominant, and produces antiferromagnetic ordering on the two diagonal sublattices; and then the effect of the J_1 interaction is to align the two sublattices to form a columnar ordered state as illustrated in Figure 2, an example of the ‘order by disorder’ phenomenon. Numerical investigations [3, 4, 5, 7, 8, 9, 10, 13] have shown that the boundaries of these two phases lie at $\alpha \simeq 0.38$ and $\alpha \simeq 0.60$, respectively.

The nature of the intermediate phase or phases remains controversial. Monte Carlo simulations are hampered by the ‘minus sign’ problem; exact diagonalizations are limited to small lattices; and series expansions are based on some particular ordered reference state, and




FIG. 1: Lattice spin models with frustration. Solid lines represent nearest-neighbour antiferromagnetic interactions J_1 ; dashed lines represent next-nearest-neighbour interactions J_2 . Case: a) anisotropic triangular lattice; b) $J_1 - J_2$ model; c) Shastry-Sutherland model; d) Union Jack model.

are only valid within a single phase. It is generally believed that the intermediate phase is gapped and shows no long-range magnetic order. Field theory approaches [5, 6] and dimer series expansions [5, 7, 8, 9] seem to indicate a columnar dimerized state in the intermediate region, with spontaneous breaking of translational symmetry, as illustrated in Figure 2. Capriotti, Sorella and coworkers, on the other hand, have suggested a homogeneous spin-liquid ‘plaquette RVB’ state [10, 11], and have found that exact diagonalization up to 6x6 sites shows no strong evidence of dimerization [12]. Another Monte Carlo study has suggested a columnar dimer state with plaquette-type modulation [13]. Sushkov *et al.* [9] have even suggested that there may be three different phases in the intermediate region: reading from left to




FIG. 2: Phase diagram of the $J_1 - J_2$ model.

right, a Néel state with columnar dimerization; a columnar dimerized spin liquid, and a columnar dimerized spin liquid with plaquette-type modulation.

Several discussions have centered on the Lieb-Schulz-Mattis theorem in higher dimensions [14, 15, 16], which shows that for a spin system with half-integer spin per unit cell, there is an excitation energy gap behaving like $1/L$, where L is the linear size of the system. Takano *et al.* [17] have argued that a uniform RVB state without gapless singlet excitations is excluded by the theorem, and that the true ground state is a plaquette state with spontaneously broken translation invariance and fourfold degeneracy. Later arguments [11, 16, 18] have refuted this, however, and shown that the theorem may be satisfied if the translation symmetry remains unbroken, but the ground state has a fourfold topological degeneracy instead, as in a simple dimer model.

The Hamiltonian for the Union Jack model is

$$H = J_1 \sum_{\langle nn \rangle} \mathbf{S}_i \cdot \mathbf{S}_j + J_2 \sum_{A: \langle nnn \rangle} \mathbf{S}_i \cdot \mathbf{S}_j \quad (1)$$

where the J_1 and J_2 interactions connect sites as shown in Figure 1d). A classical variational analysis shows that for $\alpha = J_2/J_1 < 0.5$, the ground state is the Néel state, as in the $J_1 - J_2$ model. For $\alpha > 0.5$, the ground state is the canted ferrimagnetic state shown in Figure 3, where the spins on the A sublattice are canted at an angle θ to those on the B sublattice, and 2θ to their neighbours on the A sublattice. The energy of this state is

$$E_0 = NS^2[\alpha \cos 2\theta - 2 \cos \theta] \quad (2)$$

where we have set $J_1 = 1$, S is the total spin per site, and N is the number of sites. This energy is minimized when

$$\sin \theta (2\alpha \cos \theta - 1) = 0. \quad (3)$$

For $\alpha < 0.5$, the lowest energy corresponds to $\sin \theta = 0$, i.e. the simple Néel state. For $\alpha > 0.5$, the lowest energy solution is

$$\cos \theta = \frac{1}{2\alpha}, \quad (4)$$

corresponding to the canted state. In the limit $\alpha \rightarrow \infty$, the angle $\theta \rightarrow \pi/2$: the spins on the A sublattice are




FIG. 3: Spin configurations in the canted phase.

Néel ordered, as expected, and the spins on the A and B sublattices are at right angles to each other.

In the canted phase, there is a net magnetization per site in the z direction,

$$M_z = \frac{S}{2}(1 - \cos \theta) = \frac{S}{2}\left(1 - \frac{1}{2\alpha}\right) \quad (5)$$

and a staggered magnetization per site

$$M_s = \frac{S}{2}(1 + \cos \theta) = \frac{S}{2}\left(1 + \frac{1}{2\alpha}\right). \quad (6)$$

The model thus demonstrates a new mechanism for causing ferrimagnetism. Ferrimagnetism usually arises when the individual ionic spins are different on different sublattices. Here the spins have the same magnitude on each sublattice, and the interactions are antiferromagnetic, but the frustration between them produces a nonzero overall magnetic moment.

In the neighbourhood of the transition point, $\alpha_c = 0.5$, we find for the ground state energy per bond $\epsilon_0 = E_0/2N$ in zero magnetic field :

$$\epsilon_0 \sim -\frac{3S^2}{8}, \quad \alpha \rightarrow \alpha_c \quad (7)$$

$$\frac{d\epsilon_0}{d\alpha} \sim S^2/2, \quad \alpha \rightarrow \alpha_c \quad (8)$$

$$\frac{d^2\epsilon_0}{d\alpha^2} \sim \begin{cases} 0, & \alpha \rightarrow \alpha_c^- \\ -4S^2, & \alpha \rightarrow \alpha_c^+ \end{cases} \quad (9)$$

while for the net magnetization in the z direction,

$$M_z \sim S(\alpha - \alpha_c), \quad \alpha \rightarrow \alpha_c. \quad (10)$$

Thus the transition is not quite the simple first-order transition found classically for other frustrated models. There is no “latent heat”, i.e. no discontinuity in $d\epsilon_0/d\alpha$,

although there is a finite discontinuity in the second derivative. The magnetization M_z vanishes linearly at α_c , and the energy gap is zero on both sides of the transition, corresponding to spontaneous breaking of the spin rotation symmetry. Translation symmetry on the lattice is also broken on both sides of the transition. The question then arises, whether quantum fluctuations will change the character of the transition.

In the remainder of this paper, we shall discuss how this picture is modified when one goes beyond the classical analysis to a spin-wave treatment. In Section II we shall present a modified spin-wave treatment of the Néel phase to second order; and in Section III we shall give a spin-wave treatment of the canted phase to first order. The results are discussed in Section IV. Our conclusions are summarized in Section V.

II. SPIN-WAVE THEORY FOR THE NÉEL PHASE.

A. Formulation

A review of spin-wave approaches to the $J_1 - J_2$ model has been given by Gochev [19]. Chandra and Doucot [2] gave the first conventional spin-wave treatment. The results of this approach were found to become unstable as the frustration α increases, due to strong interactions between the spin wave bosons. The interactions involve a quadratic term

$$\tilde{H}_2 = \sum_{\mathbf{k}} Q_{\mathbf{k}} (\alpha_{\mathbf{k}} \beta_{\mathbf{k}} + \alpha_{\mathbf{k}}^\dagger \beta_{\mathbf{k}}^\dagger) \quad (11)$$

which is comparable with the zeroth-order Hamiltonian for $S \sim 1$ and $\alpha \sim 1$. Thus one needs to go to a modified treatment in which this quadratic term is absent

[20, 21]: this requirement turns out to be equivalent to Takahashi's modified spin-wave theory [23]. A very similar treatment can be applied to the Union Jack model. We will follow Gochev's notation [19] as far as possible.

Let us rewrite the Hamiltonian (1), adding in a staggered magnetic field h :

$$H = \sum_{\langle lm \rangle} \mathbf{S}_l \cdot \mathbf{S}_m + \alpha \sum_{\langle l_1 l_2 \rangle} \mathbf{S}_{l_1} \cdot \mathbf{S}_{l_2} + h \left(\sum_l S_l^z - \sum_m S_m^z \right) \quad (12)$$

where we have divided the lattice into even (A) and odd (B) sublattices, denoted by indices l and m respectively, and set $J_1 = 1$.

Introduce creation and annihilation operators for the "spin deviation" on the two sublattices by means of a Dyson-Maleev transformation:

$$\begin{aligned} S_l^z &= S - a_l^\dagger a_l \\ S_l^+ &= (2S)^{1/2} a_l - (2S)^{-1/2} a_l^\dagger a_l a_l \\ S_l^- &= (2S)^{1/2} a_l^\dagger \\ S_m^z &= b_m^\dagger b_m - S \\ S_m^+ &= (2S)^{1/2} b_m^\dagger - (2S)^{-1/2} b_m^\dagger b_m^\dagger b_m \\ S_m^- &= (2S)^{1/2} b_m \end{aligned} \quad (13)$$

and then perform a Fourier transform

$$\begin{aligned} a_{\mathbf{k}} &= \left(\frac{2}{N} \right)^{1/2} \sum_l e^{i\mathbf{k} \cdot \mathbf{l}} a_l \\ b_{\mathbf{k}} &= \left(\frac{2}{N} \right)^{1/2} \sum_m e^{-i\mathbf{k} \cdot \mathbf{m}} b_m \end{aligned} \quad (14)$$

to give the Hamiltonian in the form

$$\begin{aligned} H &= -2S^2 N \left(1 - \frac{\alpha}{2} - \frac{h}{2S} \right) + 4S \left\{ \left(1 - \alpha - \frac{h}{4S} \right) \sum_{\mathbf{k}} a_{\mathbf{k}}^\dagger a_{\mathbf{k}} + \left(1 - \frac{h}{4S} \right) \sum_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{\mathbf{k}} [\gamma_{\mathbf{k}} (a_{\mathbf{k}} b_{\mathbf{k}} + a_{\mathbf{k}}^\dagger b_{\mathbf{k}}^\dagger) + \alpha \eta_{\mathbf{k}} a_{\mathbf{k}}^\dagger a_{\mathbf{k}}] \right\} \\ &\quad - \frac{4}{N} \sum_{\mathbf{1234}} \delta_{\mathbf{1-2-3+4}} [2\gamma_{\mathbf{3-4}} a_{\mathbf{1}}^\dagger a_{\mathbf{2}} b_{\mathbf{3}}^\dagger b_{\mathbf{4}} + \gamma_{\mathbf{4}} a_{\mathbf{1}}^\dagger a_{\mathbf{2}} a_{\mathbf{3}} b_{\mathbf{4}} + \gamma_{\mathbf{1}} a_{\mathbf{1}}^\dagger b_{\mathbf{2}}^\dagger b_{\mathbf{3}}^\dagger b_{\mathbf{4}}] \\ &\quad + \frac{4\alpha}{N} \sum_{\mathbf{1234}} \delta_{\mathbf{1+2-3-4}} a_{\mathbf{1}}^\dagger a_{\mathbf{2}}^\dagger a_{\mathbf{3}} a_{\mathbf{4}} (\eta_{\mathbf{2-4}} - \frac{1}{2}(\eta_{\mathbf{1}} + \eta_{\mathbf{2}})) \end{aligned} \quad (15)$$

where we have used the shorthand notation $\mathbf{1} \cdots \mathbf{4}$ for momenta $\mathbf{k}_1 \cdots \mathbf{k}_4$, and $\gamma_{\mathbf{k}}, \eta_{\mathbf{k}}$ are the structure factors for the full lattice and the A sublattice respectively:

$$\begin{aligned} \gamma_{\mathbf{k}} &= \cos \frac{k_x}{2} \cos \frac{k_y}{2} \\ \eta_{\mathbf{k}} &= \frac{1}{2} (\cos k_x + \cos k_y), \end{aligned} \quad (16)$$

if we set the spacing of each sublattice equal to 1, and k_x, k_y are the components of momentum along the diagonal axes of the two sublattices.

The Hamiltonian can now be diagonalized up to second order by a Bogoliubov transformation:

$$a_{\mathbf{k}} = u_{\mathbf{k}} \alpha_{\mathbf{k}} - v_{\mathbf{k}} \beta_{\mathbf{k}}^\dagger, \quad b_{\mathbf{k}} = u_{\mathbf{k}} \beta_{\mathbf{k}} - v_{\mathbf{k}} \alpha_{\mathbf{k}}^\dagger \quad (17)$$

where

$$u_{\mathbf{k}}^2 - v_{\mathbf{k}}^2 = 1. \quad (18)$$

After normal ordering the transformed Hamiltonian, the condition that off-diagonal quadratic terms vanish turns out to be:

$$\begin{aligned} Q_{\mathbf{k}} &= 4S[\gamma_{\mathbf{k}}(u_{\mathbf{k}}^2 + v_{\mathbf{k}}^2) - 2u_{\mathbf{k}}v_{\mathbf{k}}(1 - \frac{h}{4S} - \frac{\alpha}{2}(1 - \eta_{\mathbf{k}}))] \\ &\quad + 4(R_1 - R_2)[\gamma_{\mathbf{k}}(u_{\mathbf{k}}^2 + v_{\mathbf{k}}^2) - 2u_{\mathbf{k}}v_{\mathbf{k}}] \\ &\quad - 4\alpha u_{\mathbf{k}}v_{\mathbf{k}}(R_2 - R_3)(1 - \eta_{\mathbf{k}}) \\ &= 0 \end{aligned} \quad (19)$$

where the lattice sums R_i are

$$R_1 = \frac{2}{N} \sum_{\mathbf{k}} \gamma_{\mathbf{k}} u_{\mathbf{k}} v_{\mathbf{k}}, \quad R_2 = \frac{2}{N} \sum_{\mathbf{k}} v_{\mathbf{k}}^2, \quad R_3 = \frac{2}{N} \sum_{\mathbf{k}} \eta_{\mathbf{k}} v_{\mathbf{k}}^2 \quad (20)$$

A solution to equation (19) can easily be found (for $h \leq 0$):

$$u_{\mathbf{k}} = \left[\frac{1 + \epsilon_{\mathbf{k}}}{2\epsilon_{\mathbf{k}}} \right]^{1/2}, \quad v_{\mathbf{k}} = \text{sgn}(\gamma_{\mathbf{k}}) \left[\frac{1 - \epsilon_{\mathbf{k}}}{2\epsilon_{\mathbf{k}}} \right]^{1/2} \quad (21)$$

$$\begin{aligned} \epsilon_{\mathbf{k}} &= \left(1 - \frac{\gamma_{\mathbf{k}}^2}{f_{\mathbf{k}}^2} \right)^{1/2}, \\ f_{\mathbf{k}} &= 1 - h\sigma - \rho \frac{\alpha}{2}(1 - \eta_{\mathbf{k}}) \end{aligned} \quad (22)$$

where

$$\sigma = \frac{1}{4(S + R_1 - R_2)}$$

$$\rho = \frac{S + R_3 - R_2}{S + R_1 - R_2} \quad (23)$$

(the parameter ρ is Gochev's $\bar{\alpha}$). Then one merely has to find self-consistent solutions for the two parameters σ and ρ , given by equations (23).

We can now write the Hamiltonian as

$$H_{DM} = W_0 + H_0 + V_{DM}. \quad (24)$$

The constant term is

$$\begin{aligned} W_0 &= 2N\epsilon_0 = 2N[-(S + R_1 - R_2)^2 \\ &\quad + \frac{\alpha}{2}(S + R_3 - R_2)^2 + \frac{h}{2}(S - R_2)] \end{aligned} \quad (25)$$

(note that there is a misprint in Gochev [19] at this point). The quadratic part H_0 is diagonal:

$$H_0 = \sum_{\mathbf{k}} (E_{\mathbf{k}}^{\alpha} \alpha_{\mathbf{k}}^{\dagger} \alpha_{\mathbf{k}} + E_{\mathbf{k}}^{\beta} \beta_{\mathbf{k}}^{\dagger} \beta_{\mathbf{k}}) \quad (26)$$

where the α and β bosons have different spin-wave energies in this case:

$$\begin{aligned} E_{\mathbf{k}}^{\alpha} &= 4S[u_{\mathbf{k}}^2(1 - \alpha(1 - \eta_{\mathbf{k}}) - \frac{h}{4S}) + v_{\mathbf{k}}^2(1 - \frac{h}{4S}) \\ &\quad - 2u_{\mathbf{k}}v_{\mathbf{k}}\gamma_{\mathbf{k}}] + 4(R_1 - R_2)(u_{\mathbf{k}}^2 + v_{\mathbf{k}}^2 - 2u_{\mathbf{k}}v_{\mathbf{k}}\gamma_{\mathbf{k}}) \\ &\quad + 4\alpha u_{\mathbf{k}}^2(R_2 - R_3)(1 - \eta_{\mathbf{k}}) \end{aligned} \quad (27)$$

and $E_{\mathbf{k}}^{\beta}$ is a similar expression with $u_{\mathbf{k}}$ and $v_{\mathbf{k}}$ interchanged.

The normal-ordered quartic interaction operator V_{DM} is

$$\begin{aligned} V_{DM} &= -\frac{2}{N} \sum_{\mathbf{1234}} \delta_{\mathbf{1+2-3-4}} [\Phi^{(1)} \alpha_{\mathbf{1}} \alpha_{\mathbf{2}} \beta_{\mathbf{3}} \beta_{\mathbf{4}} + \Phi^{(2)} \alpha_{\mathbf{3}}^{\dagger} \alpha_{\mathbf{4}}^{\dagger} \beta_{\mathbf{1}}^{\dagger} \beta_{\mathbf{2}}^{\dagger} - 2\Phi^{(3)} \alpha_{\mathbf{3}}^{\dagger} \beta_{\mathbf{4}} \alpha_{\mathbf{1}} \alpha_{\mathbf{2}} - 2\Phi^{(4)} \alpha_{\mathbf{4}}^{\dagger} \beta_{\mathbf{1}}^{\dagger} \beta_{\mathbf{2}}^{\dagger} \beta_{\mathbf{3}} \\ &\quad - 2\Phi^{(5)} \beta_{\mathbf{2}}^{\dagger} \beta_{\mathbf{3}} \beta_{\mathbf{4}} \alpha_{\mathbf{1}} - 2\Phi^{(6)} \alpha_{\mathbf{3}}^{\dagger} \alpha_{\mathbf{4}}^{\dagger} \beta_{\mathbf{1}}^{\dagger} \alpha_{\mathbf{2}} + \Phi^{(7)} \beta_{\mathbf{1}}^{\dagger} \beta_{\mathbf{2}}^{\dagger} \beta_{\mathbf{3}} \beta_{\mathbf{4}} + \Phi^{(8)} \alpha_{\mathbf{3}}^{\dagger} \alpha_{\mathbf{4}}^{\dagger} \alpha_{\mathbf{1}} \alpha_{\mathbf{2}} + 4\Phi^{(9)} \alpha_{\mathbf{4}}^{\dagger} \beta_{\mathbf{1}}^{\dagger} \alpha_{\mathbf{2}} \beta_{\mathbf{3}}]. \end{aligned} \quad (28)$$

Explicit expressions for the vertex functions $\Phi^{(i)}(\mathbf{1234})$ of the Néel phase are given in Appendix A.

B. Higher-order corrections

To order 1 in a $1/S$ expansion, the ground-state energy per bond is ϵ_0 , as given by equation (25). The staggered magnetization per site is

$$M_s = 2 \frac{\partial \epsilon_0}{\partial h} \Big|_{h=0} = S - R_2, \quad (29)$$

and the spin-wave energy is that given by equation (27).

We can now use perturbation theory to calculate the next-order ($1/S$) corrections to these results. The leading correction to the ground-state energy corresponds to Figure 4a), and is given by

$$\frac{\Delta E_0}{2N} = - \left(\frac{2}{N} \right)^3 \sum_{\mathbf{1234}} \delta_{\mathbf{1+2-3-4}} \frac{\Phi^{(1)}(\mathbf{1234}) \Phi^{(2)}(\mathbf{3412})}{(E_{\mathbf{1}}^{\alpha} + E_{\mathbf{2}}^{\alpha} + E_{\mathbf{3}}^{\beta} + E_{\mathbf{4}}^{\beta})}, \quad (30)$$

while the corrections to the spin-wave energies $E_{\mathbf{k}}^{\alpha}$ and $E_{\mathbf{k}}^{\beta}$ (Fig. 4b) are

$$\begin{aligned}\Delta E_{\mathbf{k}}^{\alpha} &= -8 \left(\frac{2}{N} \right)^2 \sum_{123} \delta_{2+3-1-\mathbf{k}} \frac{\Phi^{(6)}(\mathbf{1k23})\Phi^{(3)}(\mathbf{23k1})}{(E_1^{\beta} + E_2^{\alpha} + E_3^{\alpha} - E_{\mathbf{k}}^{\alpha})} - 8 \left(\frac{2}{N} \right)^2 \sum_{123} \delta_{1+2-3-\mathbf{k}} \frac{\Phi^{(2)}(\mathbf{123k})\Phi^{(1)}(\mathbf{k321})}{(E_1^{\beta} + E_2^{\beta} + E_3^{\alpha} + E_{\mathbf{k}}^{\alpha})}, \\ \Delta E_{\mathbf{k}}^{\beta} &= -8 \left(\frac{2}{N} \right)^2 \sum_{123} \delta_{2+3-1-\mathbf{k}} \frac{\Phi^{(5)}(\mathbf{1k23})\Phi^{(4)}(\mathbf{23k1})}{(E_1^{\alpha} + E_2^{\beta} + E_3^{\beta} - E_{\mathbf{k}}^{\beta})} - 8 \left(\frac{2}{N} \right)^2 \sum_{123} \delta_{1+2-3-\mathbf{k}} \frac{\Phi^{(1)}(\mathbf{123k})\Phi^{(2)}(\mathbf{k321})}{(E_1^{\alpha} + E_2^{\alpha} + E_3^{\beta} + E_{\mathbf{k}}^{\beta})}\end{aligned}\quad (31)$$




FIG. 4: Perturbation diagrams in the Néel phase. Solid lines correspond to α bosons, dashed lines to β bosons. a) Leading correction to the ground-state energy. b) Leading corrections to the spin-wave energy $E_{\mathbf{k}}^{\alpha}$.

The corrections to the staggered magnetization can also be represented in terms of perturbation diagrams as follows. We have

$$M_s = \frac{1}{N} \langle \psi_0 | \sum_l S_l^z - \sum_m S_m^z | \psi_0 \rangle \quad (32)$$

$$= \langle \psi_0 | S - R_2 - V_M^I - V_M^{II} | \psi_0 \rangle \quad (33)$$

where

$$V_M^I = \sum_{\mathbf{k}} (u_{\mathbf{k}}^2 + v_{\mathbf{k}}^2) (\alpha_{\mathbf{k}}^{\dagger} \alpha_{\mathbf{k}} + \beta_{\mathbf{k}}^{\dagger} \beta_{\mathbf{k}}) \quad (34)$$

$$V_M^{II} = \sum_{\mathbf{k}} 2u_{\mathbf{k}}v_{\mathbf{k}} (\alpha_{\mathbf{k}}^{\dagger} \beta_{\mathbf{k}}^{\dagger} + \alpha_{\mathbf{k}} \beta_{\mathbf{k}}) \quad (35)$$

Hence the leading corrections to M_s of $O(1/S)$ correspond to the diagrams shown in Figure 5, and are given by

$$\Delta M_s = -\frac{16}{N^3} \sum_{1234} \delta_{1+2-3-4} \frac{\Phi^{(1)}(\mathbf{1234})\Phi^{(2)}(\mathbf{3412})}{(E_1^{\alpha} + E_2^{\alpha} + E_3^{\beta} + E_4^{\beta})^2} \left[\frac{1}{\epsilon_1} + \frac{1}{\epsilon_2} + \frac{1}{\epsilon_3} + \frac{1}{\epsilon_4} \right] \quad (36)$$

$$-\frac{32}{N^3} \sum_{1234} \delta_{1+2-3-4} \frac{F(\mathbf{1234})}{(E_1^{\alpha} + E_2^{\alpha} + E_3^{\beta} + E_4^{\beta})} \quad (37)$$

where

$$F(\mathbf{1234}) = \Phi^{(1)}(\mathbf{1234}) \left[\frac{\gamma_1 \Phi^{(4)}(\mathbf{3412})}{f_1 \epsilon_1 (E_1^{\alpha} + E_1^{\beta})} + \frac{\gamma_3 \Phi^{(6)}(\mathbf{4312})}{f_3 \epsilon_3 (E_3^{\alpha} + E_3^{\beta})} \right] + \Phi^{(2)}(\mathbf{3412}) \left[\frac{\gamma_1 \Phi^{(5)}(\mathbf{2134})}{f_1 \epsilon_1 (E_1^{\alpha} + E_1^{\beta})} + \frac{\gamma_3 \Phi^{(3)}(\mathbf{1234})}{f_3 \epsilon_3 (E_3^{\alpha} + E_3^{\beta})} \right]. \quad (38)$$

The results of these calculations are discussed in Section IV.

III. SPIN-WAVE THEORY FOR THE CANTED PHASE.

In this section we shall present a modified spin wave treatment of the canted phase. The formalism turns out

to be somewhat more complicated in this case, and we shall only keep terms down to $O(S)$ in the energy. We write the Hamiltonian as

$$\begin{aligned}H &= \sum_{\langle nn \rangle} \mathbf{S}_i \cdot \mathbf{S}_j + \alpha \sum_{A: \langle nnn \rangle} \mathbf{S}_i \cdot \mathbf{S}_j \\ &+ h_1 \sum_{B:i} S_i^z + h_2 \sum_{A:i} S_i^z,\end{aligned}\quad (39)$$




FIG. 5: Perturbation diagrams contributing to the Néel magnetization at order $1/S$.

$$\begin{aligned}
H = & \sum_{B:\mathbf{n},\mu} \left[-(S_{B:\mathbf{n}}^z S_{A:\mathbf{n}+\mu}^z + S_{B:\mathbf{n}}^x S_{A:\mathbf{n}+\mu}^x) \cos \theta + \eta_{\mathbf{n}\mu} (S_{B:\mathbf{n}}^z S_{A:\mathbf{n}+\mu}^x - S_{B:\mathbf{n}}^x S_{A:\mathbf{n}+\mu}^z) \sin \theta + S_{B:\mathbf{n}}^y S_{A:\mathbf{n}+\mu}^y \right] \\
& + \alpha \sum_{A1:\mathbf{n},\mu'} \left[(S_{A1:\mathbf{n}}^z S_{A2:\mathbf{n}+\mu'}^z + S_{A1:\mathbf{n}}^x S_{A2:\mathbf{n}+\mu'}^x) \cos 2\theta + S_{A1:\mathbf{n}}^y S_{A2:\mathbf{n}+\mu'}^y + (S_{A1:\mathbf{n}}^z S_{A2:\mathbf{n}+\mu'}^x - S_{A1:\mathbf{n}}^x S_{A2:\mathbf{n}+\mu'}^z) \sin 2\theta \right] \\
& + h_1 \sum_{B:\mathbf{n}} S_{B:\mathbf{n}}^z + h_2 \sum_{A:\mathbf{n}} S_{A:\mathbf{n}}^z
\end{aligned} \tag{40}$$

where the direction vectors are $\{\mu\} = \pm\mathbf{i}, \pm\mathbf{j}$; $\{\mu'\} = \pm(\mathbf{i} \pm \mathbf{j})$ and the phase factor $\eta_{\mathbf{n}\mu} = \pm(-1)^{n_y}$ for μ equals $\pm\mathbf{i}$ or $\pm\mathbf{j}$, respectively.

Now we introduce boson creation and annihilation operators for the spin deviation on the two sublattices by means of the Dyson-Maleev transformation:

$$\begin{aligned}
S_{A:\mathbf{m}}^z &= S - a_{\mathbf{m}}^\dagger a_{\mathbf{m}} \\
S_{A:\mathbf{m}}^+ &= (2S)^{1/2} a_{\mathbf{m}} \\
S_{A:\mathbf{m}}^- &= (2S)^{1/2} a_{\mathbf{m}}^\dagger - (2S)^{-1/2} a_{\mathbf{m}}^\dagger a_{\mathbf{m}}^\dagger a_{\mathbf{m}} \\
S_{B:\mathbf{n}}^z &= S - b_{\mathbf{n}}^\dagger b_{\mathbf{n}} \\
S_{B:\mathbf{n}}^+ &= (2S)^{1/2} b_{\mathbf{n}} - (2S)^{-1/2} b_{\mathbf{n}}^\dagger b_{\mathbf{n}} b_{\mathbf{n}}
\end{aligned}$$

quantizing the spins with respect to the configuration shown in Figure 3, so that the z axis on the B sublattice points upwards, and the z axes on the sublattices A_1 and A_2 are canted at angle θ to the downwards direction as shown, where θ is a parameter to be determined. The y axes are taken to lie perpendicular to the paper in each case. In terms of spin components, we then have:

$$S_{B:\mathbf{n}}^- = (2S)^{1/2} b_{\mathbf{n}}^\dagger \tag{41}$$

and perform a Fourier transform

$$\begin{aligned}
a_{\mathbf{k}} &= \left(\frac{2}{N} \right)^{1/2} \sum_{\mathbf{n}} e^{i\mathbf{k} \cdot \mathbf{n}} a_{\mathbf{n}}; \\
b_{\mathbf{k}} &= \left(\frac{2}{N} \right)^{1/2} \sum_{\mathbf{m}} e^{i\mathbf{k} \cdot \mathbf{m}} b_{\mathbf{m}}
\end{aligned} \tag{42}$$

to obtain

$$\begin{aligned}
H = & -NS^2 \left(2 \cos \theta - \alpha \cos 2\theta - \frac{(h_1 + h_2)}{2S} \right) + 2(NS^3)^{1/2} (\sin \theta - \alpha \sin 2\theta) (a_{\pi, \pi} + a_{\pi, \pi}^\dagger) \\
& + 2S \left[(2 \cos \theta - \frac{h_1}{2S}) \sum_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + (2(\cos \theta - \alpha \cos 2\theta) - \frac{h_2}{2S}) \sum_{\mathbf{k}} a_{\mathbf{k}}^\dagger a_{\mathbf{k}} - (1 + \cos \theta) \sum_{\mathbf{k}} \gamma_{\mathbf{k}} (b_{\mathbf{k}} a_{-\mathbf{k}} + b_{\mathbf{k}}^\dagger a_{-\mathbf{k}}^\dagger) \right. \\
& \left. + (1 - \cos \theta) \sum_{\mathbf{k}} \gamma_{\mathbf{k}} (b_{\mathbf{k}} a_{\mathbf{k}}^\dagger + b_{\mathbf{k}}^\dagger a_{\mathbf{k}}) + \frac{\alpha}{2} \sum_{\mathbf{k}} \eta_{\mathbf{k}} [2(1 + \cos 2\theta) a_{\mathbf{k}}^\dagger a_{\mathbf{k}} - (1 - \cos 2\theta) (a_{\mathbf{k}} a_{-\mathbf{k}} + a_{\mathbf{k}}^\dagger a_{-\mathbf{k}}^\dagger)] \right] \quad (43)
\end{aligned}$$

to order S , where as usual

$$\gamma_{\mathbf{k}} = \frac{1}{4} \sum_{\mu} e^{i\mathbf{k} \cdot \mu}, \quad \eta_{\mathbf{k}} = \frac{1}{4} \sum_{\mu'} e^{i\mathbf{k} \cdot \mu'}. \quad (44)$$

The term of order S^2 is simply the classical energy of this configuration, equation (2). The term of order $S^{3/2}$, proportional to $(b_{\pi, \pi} + b_{\pi, \pi}^\dagger)$, would indicate that we have not chosen the optimum reference state, and should be set to zero: this gives a condition on the angle θ identical to the classical condition, equation (3). At higher orders, this condition would give a ‘renormalised’ value for the angle θ .

The term of order S is quadratic in fermion operators, and can be diagonalized by a Bogoliubov transformation. In the sector involving momenta $(\pm \mathbf{k})$, the quadratic part of the Hamiltonian is

$$H_{\mathbf{k}} = 2S \left[\begin{pmatrix} a_{\mathbf{k}}^\dagger & b_{-\mathbf{k}} & b_{\mathbf{k}}^\dagger & a_{-\mathbf{k}} \end{pmatrix} \hat{H}_{\mathbf{k}} \begin{pmatrix} a_{\mathbf{k}} \\ b_{-\mathbf{k}}^\dagger \\ b_{\mathbf{k}} \\ a_{-\mathbf{k}}^\dagger \end{pmatrix} + N_{\mathbf{k}} \right] \quad (45)$$

where to leading order the matrix $\hat{H}_{\mathbf{k}}$ has elements h_{ij} given by

$$h_{11} = h_{44} = 2(\cos \theta - \alpha \cos 2\theta) + \alpha(1 + \cos 2\theta)\eta_{\mathbf{k}} - \frac{h_2}{2S}; \quad (46)$$

$$h_{22} = h_{33} = 2 \cos \theta - \frac{h_1}{2S}; \quad (47)$$

$$h_{12} = h_{21} = h_{34} = h_{43} = -(1 + \cos \theta)\gamma_{\mathbf{k}}; \quad (48)$$

$$h_{13} = h_{31} = h_{24} = h_{42} = (1 - \cos \theta)\gamma_{\mathbf{k}}; \quad (49)$$

$$h_{23} = h_{32} = 0. \quad (50)$$

$$h_{14} = h_{41} = -\alpha(1 - \cos 2\theta)\eta_{\mathbf{k}}; \quad (51)$$

Note that the matrix $\hat{H}_{\mathbf{k}}$ is symmetric about *both* diagonals. The normal-ordering correction to $O(S)$ is

$$N_{\mathbf{k}} = -h_{11} - h_{22} \quad (52)$$

Thus the procedure here involves a 4x4 matrix diagonalization, rather than 2x2. We want a transformation which preserves commutation relations, and the symmetries of the problem, i.e. we need to find a *symplectic* transformation:

$$\begin{pmatrix} a_{\mathbf{k}} \\ b_{-\mathbf{k}}^\dagger \\ b_{\mathbf{k}} \\ a_{-\mathbf{k}}^\dagger \end{pmatrix} = S_{\mathbf{k}} \begin{pmatrix} \alpha_{\mathbf{k}} \\ \beta_{-\mathbf{k}}^\dagger \\ \beta_{\mathbf{k}} \\ \alpha_{-\mathbf{k}}^\dagger \end{pmatrix} \quad (53)$$

where the elements s_{ij} of $S_{\mathbf{k}}$ obey

$$\begin{aligned}
\sum_j s_{ij}^2 (-1)^{j+1} &= (-1)^{i+1} \\
\sum_j s_{ij} s_{kj} (-1)^{j+1} &= 0 \\
s_{ij} &= s_{5-i, 5-j} \quad (54)
\end{aligned}$$

Then

$$\hat{H}'_{\mathbf{k}} = S^T \hat{H}_{\mathbf{k}} S. \quad (55)$$

There are initially 16 unknown parameters, corresponding to the elements of the transformation matrix $S_{\mathbf{k}}$. The conditions (54) turn out to eliminate 12 of these, leaving only 4 independent parameters; these 4 parameters are determined by the condition that $\hat{H}'_{\mathbf{k}}$ should be diagonal, i.e.

$$h'_{12} = h'_{13} = h'_{14} = h'_{23} = 0. \quad (56)$$

We have determined the solution to this problem numerically. In its diagonalized form, the Hamiltonian now reads:

$$\begin{aligned}
H_{\mathbf{k}} = & 2S [h'_{11} (a_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} + \alpha_{-\mathbf{k}}^\dagger \alpha_{-\mathbf{k}}) \\
& + h'_{22} (\beta_{\mathbf{k}}^\dagger \beta_{\mathbf{k}} + \beta_{-\mathbf{k}}^\dagger \beta_{-\mathbf{k}}) + N'_{\mathbf{k}}], \quad (57)
\end{aligned}$$

where

$$N'_{\mathbf{k}} = h'_{11} + h'_{22} - h_{11} - h_{22} \quad (58)$$

Hence one can easily obtain numerical results for the ground state energy and single-particle dispersion relations. The staggered magnetization is given by

$$M_s = \frac{1}{N} \langle \psi_0 | \sum_B S_B^z + \cos \theta \sum_A S_A^z | \psi_0 \rangle$$

TABLE I: Ground-state energy per bond as a function of α . Linear spin-wave theory: $\epsilon_0^{(1)}$; modified second-order spin-wave theory: $\epsilon_0^{(2)}$; with higher-order corrections $\epsilon_0^{(3)}$.

α	Néel expansion			Canted expansion
	$\epsilon_0^{(1)}$	$\epsilon_0^{(2)}$	$\epsilon_0^{(3)}$	
0.0	-0.32897	-0.33521	-0.33503	
0.1	-0.31778	-0.32523	-0.32496	
0.2	-0.30666	-0.31543	-0.31509	
0.3	-0.29561	-0.30586	-0.30543	
0.4	-0.28465	-0.29652	-0.29602	
0.5	-0.27377	-0.28746	-0.28691	-0.27377
0.6	-0.26300	-0.27870	-0.27817	-0.25465
0.7	-0.25233	-0.27028	-0.26990	-0.24458
0.8	-0.24180	-0.26224	-0.26224	-0.24050
0.9	-0.23141	-0.25462	-	-0.24058
1.0	-0.22119	-0.24746	-	-0.24367

TABLE II: Net magnetization M_z and staggered magnetization M_s as functions of α . Conventions as for Table I.

α	Néel expansion			Canted expansion	
	$M_s^{(1)}$	$M_s^{(2)}$	$M_s^{(3)}$	$M_s^{(1)}$	$M_z^{(1)}$
0.0	0.30340	0.30340	0.30601		
0.1	0.29582	0.29149	0.29499		
0.2	0.28759	0.27819	0.28265		
0.3	0.27859	0.26327	0.26872		
0.4	0.26873	0.24647	0.25270		
0.5	0.25785	0.22751	0.23386	0.25785	0.00000
0.6	0.24578	0.20607	0.21088	0.28800	0.02688
0.7	0.23230	0.18181	0.18163	0.30067	0.05143
0.8	0.21714	0.15442	0.14529	0.30525	0.07207
0.9	0.19992	0.12369	-	0.30579	0.08912
1.0	0.18017	0.08954	-	0.30429	0.10321

$$= \frac{S}{2}(1 + \cos \theta) - \frac{2}{N} \sum_k [s_{21}^2 + s_{23}^2 + \cos \theta (S_{12}^2 + s_{14}^2)] \quad (59)$$

and the net magnetization in the z direction:

$$M_z = \frac{1}{N} \langle \psi_0 | \sum_B S_B^z - \cos \theta \sum_A S_A^z | \psi_0 \rangle = \frac{S}{2}(1 - \cos \theta) - \frac{2}{N} \sum_k [s_{21}^2 + s_{23}^2 - \cos \theta (S_{12}^2 + s_{14}^2)] \quad (60)$$




FIG. 6: Ground-state energy per bond as a function of coupling α . Short-dashed line: classical result; long-dashed line: linear spin-wave theory; solid line: 2nd order spin-wave theory.

IV. NUMERICAL RESULTS

Numerical results for the model have been obtained using the finite-lattice method. The momentum sums were carried out for a fixed sublattice size $L = \sqrt{N}/2$, using discrete values for the momenta k_x and k_y , e.g.

$$k_x(i) = \frac{2\pi(i - 1/2)}{L}, \quad i = 1, \dots, L, \quad (61)$$

where we take half-integer values corresponding to anti-periodic boundary conditions to avoid any integrable singularities at $\mathbf{k} = 0$. Results were obtained for $L = 8, 9 \dots 12$, and a fit in powers of $1/L$ was made to extrapolate to the bulk limit $L \rightarrow \infty$. The finite-size corrections for the ground-state energy per bond scale asymptotically like $1/L^3$, and those for the magnetization like $1/L$ [24, 25]. The resulting bulk estimates are shown in Tables I and II. The values at $\alpha = 0$ agree with those obtained by Gochev [19] for the pure Heisenberg case.

Figure 6 shows the behaviour of the ground-state energy resulting from the classical calculation and linear spin-wave theory for both the Néel phase and the canted phase, and the modified treatment with corrections for the Néel phase. In linear spin-wave theory, the Néel and canted results coincide at $\alpha = 0.5$, as they must do since $\theta = 0$ there. Somewhat surprisingly, however, the Néel energy remains the lower of the two beyond that point, until the two curves cross once more at $\alpha \simeq 0.84$. In other words, the transition between the Néel and canted phases is pushed out to $\alpha \simeq 0.84$ in linear spin-wave theory, and is clearly first-order. The modified treatment to second order lowers the energy a little further, while the higher-order corrections are virtually negligible, and indistinguishable on the diagram.

Figure 7 shows the magnetizations M_s and M_z as func-




FIG. 7: Staggered magnetization M_s and net magnetization M_z as functions of α .

tions of the coupling α . In linear spin-wave theory, the staggered magnetization M_s in the Néel phase is reduced by quantum fluctuations, as expected, but the effective coupling is only half that in the $J_1 - J_2$ model, and so M_s remains substantial at $\alpha = 0.5$. In the modified second-order treatment it is lowered somewhat further, while the higher-order corrections are small and positive for low α , and turn negative beyond $\alpha \simeq 0.7$. The staggered magnetization in the canted phase is also reduced by quantum fluctuations, although the effect is reduced at large α .

Figure 8 illustrates the spin-wave dispersion of the α and β bosons in the Néel phase as given by the second-order theory. It can be seen that the dispersion curve for the β bosons remains virtually unchanged at all α . That for the α bosons, however, develops an instability at $\mathbf{k} = (\pi, \pi)$, and the energy gap is predicted to vanish when α gets too large, signalling a transition. The instability occurs at $\alpha = 1.0$ in linear spin wave theory, and $\alpha = 0.645$ in the modified second-order theory.

Figure 9 shows similar plots from the linear theory in the canted phase. The energy gap for the β bosons remains zero at all couplings, with linear dispersion near the origin; but as soon as we move away from $\alpha = 0.5$ (i.e. $\theta = 0$) an energy gap opens up for the α bosons, with quadratic dispersion near the origin. This is in accordance with the general counting of Goldstone modes in non-relativistic systems [26], and with previous discussions of Goldstone modes in canted phases [27, 28, 29], where the spontaneous symmetry breaking pattern is $SU(2) \rightarrow Z_2$.

V. CONCLUSIONS

The Union Jack model is another example of a spin-1/2 Heisenberg antiferromagnet with frustration on the square lattice. A classical variational analysis predicts a transition at $\alpha = 0.5$ between the Néel phase at small




FIG. 8: Dispersion curves in the Néel phase along high-symmetry lines in the Brillouin zone for various α . Curves a) and b) are from linear spin-wave theory, c) and d) from the modified second-order theory. Cases a) and c): α bosons; cases b) and d): β bosons.




FIG. 9: Dispersion curves in the canted phase along high-symmetry lines in the Brillouin zone, obtained from linear spin-wave theory. Case a): α bosons; b) β bosons.

α , and a canted ferrimagnetic phase at large α . Thus the system exhibits a new phenomenon, namely ferrimagnetism induced by frustration.

In linear spin-wave theory, the transition is pushed out to $\alpha_c \simeq 0.84$. This makes the model potentially much more interesting, because it is very likely that the Néel magnetization will vanish at or before that point. In the $J_1 - J_2$ model, numerical studies [3, 4, 5, 7, 8, 9, 10, 13] show that the Néel magnetization vanishes at about $\alpha_c \simeq 0.38$. The frustration in the Union Jack model is only half as strong, as seen by a glance at Figure 1 or from linear spin-wave theory, and so one might expect the magnetization to vanish at about $\alpha_c \simeq 0.76$ in

this model. A plausible scenario, then, is that a second-order Néel transition might occur at $\alpha_c \simeq 0.76$, possibly followed by an intermediate spin-liquid phase as in the $J_1 - J_2$ model, and then a first-order transition to the canted phase at somewhat larger α . Numerical experiments would be necessary to ascertain if this is indeed the case.

Acknowledgments

We would like to thank Profs. J. Oitmaa and O. Sushkov, and Dr. G. Misguich, for very useful discus-

sions and advice. This work forms part of a research project supported by a grant from the Australian Research Council.

APPENDIX

The vertex functions $\Phi^{(i)}, i = 1 \cdots 9$ are:

$$\begin{aligned} \Phi^{(1)}(\mathbf{1234}) = & \gamma(4-1)v_1u_2v_3u_4 + \gamma(4-2)u_1v_2v_3u_4 + \gamma(3-1)v_1u_2u_3v_4 + \gamma(3-2)u_1v_2u_3v_4 \\ & - \gamma(4)u_1u_2v_3u_4 - \gamma(3)u_1u_2u_3v_4 - \gamma(3)v_1v_2v_3u_4 - \gamma(4)v_1v_2u_3v_4 \\ & - \alpha Qu_1u_2v_3v_4 \end{aligned} \quad (62)$$

$$\begin{aligned} \Phi^{(2)}(\mathbf{1234}) = & \gamma(4-2)v_1u_2u_3v_4 + \gamma(3-2)v_1u_2v_3u_4 + \gamma(4-1)u_1v_2u_3v_4 + \gamma(3-1)u_1v_2v_3u_4 \\ & - \gamma(3)v_1v_2v_3u_4 - \gamma(4)v_1v_2u_3v_4 - \gamma(3)u_1u_2u_3v_4 - \gamma(4)u_1u_2v_3u_4 \\ & - \alpha Qv_1v_2u_3u_4 \end{aligned} \quad (63)$$

$$\begin{aligned} \Phi^{(3)}(\mathbf{1234}) = & \gamma(4-1)v_1u_2u_3u_4 + \gamma(3-1)v_1u_2v_3v_4 + \gamma(4-2)u_1v_2u_3u_4 + \gamma(3-2)u_1v_2v_3v_4 \\ & - \gamma(4)u_1u_2u_3u_4 - \gamma(3)u_1u_2v_3v_4 - \gamma(3)v_1v_2u_3u_4 - \gamma(4)v_1v_2v_3v_4 \\ & - \alpha Qu_1u_2u_3v_4 \end{aligned} \quad (64)$$

$$\begin{aligned} \Phi^{(4)}(\mathbf{1234}) = & \gamma(3-1)u_1v_2u_3u_4 + \gamma(4-1)u_1v_2v_3v_4 + \gamma(3-2)v_1u_2u_3u_4 + \gamma(4-2)v_1u_2v_3v_4 \\ & - \gamma(3)v_1v_2u_3u_4 - \gamma(4)v_1v_2v_3v_4 - \gamma(3)u_1u_2v_3v_4 - \gamma(4)u_1u_2u_3u_4 \\ & - \alpha Qv_1v_2v_3u_4 \end{aligned} \quad (65)$$

$$\begin{aligned} \Phi^{(5)}(\mathbf{1234}) = & \gamma(4-1)v_1v_2v_3u_4 + \gamma(4-2)u_1u_2v_3u_4 + \gamma(3-1)v_1v_2u_3v_4 + \gamma(3-2)u_1u_2u_3v_4 \\ & - \gamma(4)u_1v_2v_3u_4 - \gamma(3)u_1v_2u_3v_4 - \gamma(3)v_1u_2v_3u_4 - \gamma(4)v_1u_2u_3v_4 \\ & - \alpha Qu_1v_2v_3v_4 \end{aligned} \quad (66)$$

$$\begin{aligned} \Phi^{(6)}(\mathbf{1234}) = & \gamma(4-1)u_1u_2u_3v_4 + \gamma(4-2)v_1v_2u_3v_4 + \gamma(3-1)u_1u_2v_3u_4 + \gamma(3-2)v_1v_2v_3u_4 \\ & - \gamma(4)v_1u_2u_3v_4 - \gamma(3)v_1u_2v_3u_4 - \gamma(3)u_1v_2u_3v_4 - \gamma(4)u_1v_2v_3u_4 \\ & - \alpha Qv_1u_2u_3u_4 \end{aligned} \quad (67)$$

$$\begin{aligned} \Phi^{(7)}(\mathbf{1234}) = & \gamma(4-2)v_1u_2v_3u_4 + \gamma(4-1)u_1v_2v_3u_4 + \gamma(3-2)v_1u_2u_3v_4 + \gamma(3-1)u_1v_2u_3v_4 \\ & - \gamma(4)v_1v_2v_3u_4 - \gamma(3)v_1v_2u_3v_4 - \gamma(3)u_1u_2v_3u_4 - \gamma(4)u_1u_2u_3v_4 \\ & - \alpha Qv_1v_2v_3v_4 \end{aligned} \quad (68)$$

$$\begin{aligned} \Phi^{(8)}(\mathbf{1234}) = & \gamma(4-2)u_1v_2u_3v_4 + \gamma(4-1)v_1u_2u_3v_4 + \gamma(3-2)u_1v_2v_3u_4 + \gamma(3-1)v_1u_2v_3u_4 \\ & - \gamma(4)u_1u_2u_3v_4 - \gamma(3)u_1u_2v_3u_4 - \gamma(3)v_1v_2u_3v_4 - \gamma(4)v_1v_2v_3u_4 \\ & - \alpha Qu_1u_2u_3u_4 \end{aligned} \quad (69)$$

$$\begin{aligned} \Phi^{(9)}(\mathbf{1234}) = & \gamma(4-2)u_1u_2u_3u_4 + \gamma(4-1)v_1v_2u_3u_4 + \gamma(3-2)u_1u_2v_3v_4 + \gamma(3-1)v_1v_2v_3v_4 \\ & - \gamma(4)u_1v_2u_3u_4 - \gamma(3)u_1v_2v_3v_4 - \gamma(3)v_1u_2u_3u_4 - \gamma(4)v_1u_2v_3v_4 \\ & - \alpha Qv_1u_2v_3u_4 \end{aligned} \quad (70)$$

where

$$Q = \eta(3-2) + \eta(4-2) - \eta(3) - \eta(4). \quad (71)$$

-
- [1] P.W. Anderson, Science (Washington, DC, U.S.) **235**, 1196 (1987).
 - [2] P. Chandra and B. Doucot, Phys. Rev. B **38**, 9335 (1988).
 - [3] E. Dagotto and A. Moreo, Phys. Rev. Lett. **63**, 2148 (1989); F. Figueirido *et al.*, Phys. Rev. B **41**, 4619 (1990).
 - [4] H.J. Schulz, T.A.L. Ziman and D. Poilblanc, J. Phys. I **6**, 675 (1996).
 - [5] V.N. Kotov, J. Oitmaa, O.P. Sushkov and Zheng W-H., Phys. Rev. B **60**, 14613 (1999).
 - [6] N. Read and S. Sachdev, Phys. Rev. Lett. **66**, 1773 (1991); *ibid* **62**, 1694 (1989); G. Murthy and S. Sachdev, Nucl. Phys. B **344**, 557 (1990).
 - [7] M.P. Gelfand, R.R.P. Singh and D.A. Huse, Phys. Rev. B **40**, 10801 (1989); M.P. Gelfand, *ibid*, **42**, 8206 (1990).
 - [8] R.R.P. Singh, Zheng W-H., C.J. Hamer and J. Oitmaa, Phys. Rev. B **60**, 7278 (1999).
 - [9] O.P. Sushkov, J. Oitmaa and Zheng W-H., Phys. Rev. B **63**, 104420 (2001).
 - [10] L. Capriotti and S. Sorella, Phys. Rev. Lett. **84**, 3173 (2000).
 - [11] S. Sorella, L. Capriotti, F. Becca and A. Parola, Phys. Rev. Lett. **91**, 257005 (2003).
 - [12] L. Capriotti, F. Becca, A. Parola and S. Sorella, Phys. Rev. B **67**, 212402 (2003).
 - [13] M.S.L. du Croo de Jongh, J.M.J. van Leeuwen and W. van Saarloos, Phys. Rev. B **62**, 14844 (2000).
 - [14] M. Yamanaka, M. Oshikawa and I. Affleck, Phys. Rev. Lett. **79**, 1110 (1997).
 - [15] M. Oshikawa, Phys. Rev. Lett. **84**, 1535 (2000).
 - [16] M.B. Hastings, Phys. Rev. B **69**, 104431 (2004).
 - [17] K. Takano, Y. Kito, Y. Ono and K. Sano, Phys. Rev. Lett. **91**, 197202 (2003).
 - [18] G. Misguich, C. Lhuillier, M. Mambrini and P. Sindzingre, Eur. Phys. J. B **26**, 167 (2002).
 - [19] I.G. Gochev, Phys. Rev. B **49**, 9594 (1994).
 - [20] J.H. Xu and C.S. Ting, Phys. Rev. B **42**, 6861 (1990).
 - [21] H. Nishimori and Y. Saika, J. Phys. Soc. Jpn. **59**, 4454 (1990); A.F. Baranov and O.A. Starykh, JETP Lett. **51**, 311 (1990); T. Oguchi and H. Kitakani, J. Phys. Soc. Jpn. **59**, 3322 (1990).
 - [22] J.E. Hirsch and S. Tang, Phys. Rev. B **39**, 2887 (1989).
 - [23] M. Takahashi, Phys. Rev. Lett. **58**, 168 (1987); Phys. Rev. B **40**, 2494 (1989).
 - [24] P. Hasenfratz and F. Niedermayer, Z. Phys. B **92**, 91 (1993).
 - [25] Zheng W-H. and C.J. Hamer, Phys. Rev. B **47**, 7961 (1993).
 - [26] H.B. Nielsen and S. Chada, Nucl. Phys. B **105**, 445 (1976).
 - [27] S. Sachdev and T. Senthil, Ann. Phys. (N.Y.) **251**, 76 (1996).
 - [28] S. Das Sarma, S. Sachdev, and L. Zheng, Phys. Rev. B **58**, 4672 (1998).
 - [29] J.M. Roman and J. Soto, Phys. Rev. B **62**, 3300 (2000).